



HP DISINFECTANT CLEANER

Ready to Use Disinfectant Cleaner and Deodorant

ACTIVE INGREDIENTS

Hydrogen Peroxide	4.25%
OTHER INGREDIENT	95.75%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID:
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **IF SWALLOWED:** Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR TREATMENT ADVICE. Call Chem-Tel at 1-800-255-3924. Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.



HP DISINFECTANT CLEANER

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMAN AND DOMESTIC ANIMALS
CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eye or clothing. Wear chemical splash-proof goggles or face shields. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse this container to hold materials other than pesticides or diluted pesticide rinsate. Triple rinse as follows: Fill container 1/4 full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat procedure two more times. Then offer for recycling or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

This product is a one-step hospital-use germicidal cleaner and deodorant designed for general cleaning, and disinfecting, of hard, non-porous surfaces. May be used to clean and disinfect floor areas, sinks, faucets, bathrooms and tubs.

Bactericidal Activity:

Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella enterica, Escherichia coli (0157:H7), Streptococcus pneumoniae.

Antibiotic-Resistant Bactericidal Activity:

Staphylococcus aureus, Methicillin resistant (MRSA), *Staphylococcus aureus*, Community Associated Methicillin-resistant (CA-MRSA), (Genotype USA300), *Staphylococcus aureus*, Community Associated Methicillin-resistant (CA-MRSA), (Genotype USA400), *Staphylococcus epidermidis*, Methicillin-resistant (MRSE), *Streptococcus pneumoniae*, Penicillin-resistant (PRSP), *Enterococcus faecium* (Vancomycin Resistant (VRE)).

*Virucidal Activity:

*Adenovirus type 8 (5 minute contact time), *Hepatitis B Virus (HBV) (Duck Hepatitis B Virus as the surrogate) (5 minute contact time on pre-cleaned surfaces), *Hepatitis C Virus (HCV) (Bovine Viral Diarrhea Virus as the surrogate) (5 minute contact time) * Herpes Simplex Virus Type 1 (5 minute contact time), *Herpes Simplex Virus Type 2 (5 minute contact time), *HIV-1 (Human Immunodeficiency Virus Type 1) (AIDS virus) (1 minute contact time), *Influenza Virus Type A (H1N1) (1 minute contact time), *Norovirus (Feline Calicivirus as the surrogate) (5 minute contact time), *Rhinovirus Type 37 (5 minute contact time), *Rotavirus (5 minute contact time).

H1N1 Information: * This product has demonstrated effectiveness against Influenza A virus in 60 seconds and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

Modes of Application: This product can be applied by mop, sponge, cloth, paper towel, coarse trigger sprayer, or auto-scrubber. Change cloth, sponges or towels frequently to avoid redeposition of soil.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

Not recommended for use on copper, brass, granite, marble or zinc.

For Use as Ready to Use Cleaner/Disinfectant: Pre-clean visibly soiled surfaces. Apply Use Solution by coarse trigger sprayer to hard, non-porous surfaces. Spray 6 - 8 inches from surface, making sure to wet surfaces thoroughly. All surfaces must remain wet for 10 minutes. Wipe surfaces or let air dry.

NOTE: Rinsing is not necessary unless floors are to be coated with finish or restorer. When cleaning floors position "wet floor" signs around area to be cleaned. Floors will be slippery when wet or contaminated with foreign materials. Do not use on glassware, utensils, or dishes.

For use as a Virucide: Pre-clean visibly soiled surfaces.

Apply Use Solution until thoroughly wet. Let stand for 5 minutes. Wipe surfaces or let air dry.

This product kills HIV-1 (associated with AIDS), HBV and HCV on pre-cleaned surfaces/object previously soiled with blood and body fluids.

On pre-cleaned surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HBV, HCV and HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures. **Cleaning Procedures:** Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **Contact Time:** Allow surface to remain wet for 1 minute to kill HIV-1, 5 minutes to kill HBV & HCV. **Disposal of Infectious Material:** Blood and other body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

To Kill Fungi: Pre-clean visibly soiled areas. Apply use solution to hard, non-porous surfaces. Allow surface to remain wet for 10 minutes. Wipe surfaces or let air dry.

For Treatment of Animal Housing Facilities: Remove all animals and feeds from areas being treated. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals. Empty or cover all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap and detergent and rinse with water. Apply fresh Use Solution to floors, walls, cages and other washable hard, non-porous surfaces. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. To disinfect, all surfaces must remain wet for ten minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings and other closed spaces. Do not house animals or re-employ equipment until product has dried. For disinfection of feed racks, troughs, automatic feeders, fountains and watering appliances scrub with use-solution, let stand ten minutes. Then thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.

To Sanitize Non-Food Contact Surfaces: Pre-clean visibly soiled hard non-porous surfaces. Apply Use Solution until thoroughly wet. Let stand for five minutes. Wipes surfaces or let air dry. Not for use on food contact surfaces on food preparation areas.

NOTE: Do not use on glassware, utensils, or dishes.

For Use as a Deodorizer: Apply to hard, non-porous surfaces. Let stand for 10 minutes. Wipe surfaces or let air dry.

For Use as a Glass Cleaner: Apply evenly to glass surfaces through a coarse trigger spray device. Do not breathe spray. Wipe dry with an absorbent cloth or towel.

EPA REG. NO. 45745-11-14994 EPA EST. NO. 45745-TN-2

Distributed By: DURAFOG

611 Rock Spring Rd, Escondido CA 92025
(760)223-7110 | dura-fog.com
support@dura-fog.com

NET CONTENTS
5 GALS
(640 FL OZ)
(18.92L)

DISTRIBUTED BY: DURAFOG
611 ROCK SPRING RD, ESCONDIDO CA 92025 | 760-223-7110

NET CONTENTS:
5 GALS (640 FL OZ) (18.92L)